

CHAPTER 16

Solubility of organic compounds in scCO₂

Noor U Din Reshi^a, Masood Ahmad Rizvi^a, Syed Kazim Moosvi^b, Mudasir Ahmad^c, Adil Gani^c

^aDepartment of Chemistry, University of Kashmir, Srinagar, India

^bSchool Education Department, J&K Government, Srinagar, India

^cDepartment of Food Science and Technology, University of Kashmir, Srinagar, India

Contents

1. Suitability of CO ₂ as solvent for SCF	381
2. Enhancing solubility in scCO ₂	383
3. Use of models to correlate experimental solubility data	384
4. Case studies of solubility of organic compounds in scCO ₂	385
5. Amoxicillin	385
6. Antiinflammatory drugs	385
7. Artemisinin	385
8. Anesthetics	386
9. Cholesterol	386
10. Dexamethasone	386
11. Flurbiprofen	387
12. Isoniazid	387
13. 2,2'-Bipyridine and 4,4'-dimethyl-2,2'-bipyridine	387
14. Amide compounds	387
15. Polynuclear aromatic hydrocarbons	388
16. Maleic acid	388
17. Menthol	388
18. Methyl salicylate	389
19. Phenacetin	389
20. Oxymatrine	389
21. Polyacrylamide	389
22. 1,4-Dimethoxybenzene	390
23. Troeger's base	390
24. Palmitic acid + capsaicin	390
25. Phenol and pyrocatechol	390
26. 2H-chromene derivatives	391
27. Fat-soluble vitamins A, D, E, and K	391
28. Disperse dyes	392
29. Anthracene, phenanthrene, and carbazole mixture	392
30. 6-Caprolactam	392
31. Solubility of energetic materials in SCFs	393