




# Effect of hydroxy groups on structural, electronic, and biological properties of methyl carbazate containing hydroxy benzaldehyde-based imines. A combined experimental and theoretical investigation

Muzzaffar A. Bhat <sup>a</sup>  , Sajad A. Bhat <sup>a</sup> , Rusul Alabada <sup>b</sup>, Mandeep Kaur <sup>c,d</sup>,  
Harpreet Kaur <sup>e,f</sup>, G.V. Shiva Prasad <sup>g</sup>, Rajni Verma <sup>h</sup>, Anoop Dev <sup>i</sup>, Mohit Gupta <sup>j</sup>, Aashna Sinha <sup>k</sup>,  
Kholood A. Dahlous <sup>l</sup>, Raymond J. Butcher <sup>m</sup>

<sup>a</sup> Department of Chemistry, Islamic University of Science and Technology, Awantipora  
Kashmir, 192122, India

<sup>b</sup> Al-Muthanna University, College of Pharmacy, Muthanna State, Samawah 66001, Iraq

<sup>c</sup> Department of Chemistry, Jain (Deemed-to-be) University, Bengaluru, Karnataka, 560069,  
India

<sup>d</sup> Department of Sciences, Vivekananda Global University Jaipur, Rajasthan, 303012, India

<sup>e</sup> School of Basic and Applied Sciences, Shobhit University, Ganaoh, Uttar Pradesh, 247341,