



## OPEN ACCESS

### Edited by:

Mingli Liu, Morehouse School of Medicine, United States

#### Reviewed by:

Yumin Wang, Central South University, China Juni Sarkar, University of Southern California, Los Angeles, United States

#### \*Correspondence:

Mohammad Haris mharis@sidra.org Ikhlak Ahmed iahmed2@sidra.org

<sup>†</sup>These authors have contributed equally to this work and share first authorship

> <sup>‡</sup>These authors share senior authorship

#### Specialty section:

This article was submitted to Molecular Medicine, a section of the journal Frontiers in Cell and Developmental Biology

> Received: 14 October 2020 Accepted: 04 January 2021 Published: 05 February 2021

#### Citation:

Nisar S, Bhat AA, Singh M, Karedath T, Rizwan A, Hashem S, Bagga P, Reddy R, Jamal F, Uddin S, Chand G, Bedognetti D, El-Rifai W, Frenneaux MP, Macha MA, Ahmed I and Haris M (2021) Insights Into the Role of CircRNAs: Biogenesis, Characterization, Functional, and Clinical Impact in Human Malignancies. Front. Cell Dev. Biol. 9:617281. doi: 10.3389/fcell.2021.617281

# Insights Into the Role of CircRNAs: Biogenesis, Characterization, Functional, and Clinical Impact in Human Malignancies

Sabah Nisar<sup>1†</sup>, Ajaz A. Bhat<sup>1†</sup>, Mayank Singh<sup>2†</sup>, Thasni Karedath<sup>3†</sup>, Arshi Rizwan<sup>4</sup>, Sheema Hashem<sup>1</sup>, Puneet Bagga<sup>5</sup>, Ravinder Reddy<sup>6</sup>, Farrukh Jamal<sup>7</sup>, Shahab Uddin<sup>8</sup>, Gyan Chand<sup>9</sup>, Davide Bedognetti<sup>10,11,12</sup>, Wael El-Rifai<sup>13</sup>, Michael P. Frenneaux<sup>14</sup>, Muzafar A. Macha<sup>15</sup>, Ikhlak Ahmed<sup>3,16\*‡</sup> and Mohammad Haris<sup>1,17\*‡</sup>

<sup>1</sup> Functional and Molecular Imaging Laboratory, Cancer Research Department, Sidra Medicine, Doha, Qatar, <sup>2</sup> Dr. B. R. Ambedkar Institute Rotary Cancer Hospital (BRAIRCH), All India Institute of Medical Sciences (AIIMS), New Delhi, India, <sup>3</sup> Research Branch, Sidra Medicine, Doha, Qatar, <sup>4</sup> Department of Nephrology, All India Institute of Medical Sciences (AIIMS), New Delhi, India, <sup>5</sup> Diagnostic Imaging, St. Jude Children's Research Hospital, Memphis, TN, United States, <sup>6</sup> Department of Radiology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, United States, <sup>7</sup> Dr. Rammanohar Lohia Avadh University, Ayodhya, India, <sup>8</sup> Translational Research Institute, Academic Health System, Hamad Medical Corporation, Doha, Qatar, <sup>9</sup> Department of Endocrine Surgery, Sanjay Gandhi Post Graduate Institute of Medicine, Doha, Qatar, <sup>10</sup> Laboratory of Cancer Immunogenomics, Cancer Research Department, Sidra Medicine, Doha, Qatar, <sup>11</sup> Department of Internal Medicine and Medical Specialties, University of Genoa, Genoa, Italy, <sup>12</sup> College of Health and Life Sciences, Hamad Bin Khalifa University, Doha, Qatar, <sup>13</sup> Department of Surgery, University of Miami Miller School of Medicine, Miami, FL, United States, <sup>14</sup> Academic Health System, Hamad Medical Corporation, Doha, Qatar, <sup>15</sup> Watson–Crick Centre for Molecular Medicine, Islamic University of Science and Technology (IUST), Pulwama, India, <sup>16</sup> Center for Interdisciplinary Research and Innovations, University of Kashmir, Srinagar, India, <sup>17</sup> Laboratory Animal Research Center, Qatar University, Doha, Qatar

Circular RNAs (circRNAs) are an evolutionarily conserved novel class of non-coding endogenous RNAs (ncRNAs) found in the eukaryotic transcriptome, originally believed to be aberrant RNA splicing by-products with decreased functionality. However, recent advances in high-throughput genomic technology have allowed circRNAs to be characterized in detail and revealed their role in controlling various biological and molecular processes, the most essential being gene regulation. Because of the structural stability, high expression, availability of microRNA (miRNA) binding sites and tissue-specific expression, circRNAs have become hot topic of research in RNA biology. Compared to the linear RNA, circRNAs are produced differentially by backsplicing exons or lariat introns from a pre-messenger RNA (mRNA) forming a covalently closed loop structure missing 3' poly-(A) tail or 5' cap, rendering them immune to exonucleasemediated degradation. Emerging research has identified multifaceted roles of circRNAs as miRNA and RNA binding protein (RBP) sponges and transcription, translation, and splicing event regulators. CircRNAs have been involved in many human illnesses, including cancer and neurodegenerative disorders such as Alzheimer's and Parkinson's disease, due to their aberrant expression in different pathological conditions. The

1